

Statistical methods for archaeological data analysis I: Basic methods

03 - Explorative statistics & graphical display

Martin Hinz

Institut für Archäologische Wissenschaften, Universität Bern

04.03.2025

Loading data for the following steps

download data

- [muensingen_fib.csv](#)

Read the Data on Muensingen Fibulae

```
muensingen <- read.csv2("https://raw.githubusercontent.com/BernCoDALab/smada/refs/
head(muensingen)
```

```
##      X Grave Mno FL BH BFA FA CD BRA ED FEL  C   BW  BT FEW Coils Length
## 1  1   121 348 28 17  1 10 10  2  8  6 20  2.5 2.6 2.2    4    53
## 2  2   130 545 29 15  3  8  6  3  6 10 17 11.7 3.9 6.4    6    47
## 3  3   130 549 22 15  3  8  7  3 13  1 17  5.0 4.6 2.5   10    47
## 4  8   157  85 23 13  3  8  6  2 10  7 15  5.2 2.7 5.4   12    41
## 5 11   181 212 94 15  7 10 12  5 11 31 50  4.3 4.3  NA    6   128
## 6 12   193 611 68 18  7  9  9  7  3 50 18  9.3 6.5  NA    4   110
##      fibula_scheme
## 1                   B
## 2                   B
## 3                   B
## 4                   B
## 5                   C
## 6                   C
```

Cross tables (contingency tables)

For summary of data:

```
my_table <- table(muensingen$fibula_scheme, muensingen$Grave)
my_table
```

```
##
##      6 23 31 44 48 49 61 68 80 91 121 130 157 181 193
## A  1  1  1  1  0  0  0  0  0  0  0  0  0  0  0
## B  0  0  0  0  1  1  2  1  1  1  1  2  1  0  0
## C  0  0  0  0  0  0  0  0  0  0  0  0  0  1  1
```

```
addmargins(my_table)
```

```
##
##      6 23 31 44 48 49 61 68 80 91 121 130 157 181 193 Sum
## A    1  1  1  1  0  0  0  0  0  0  0  0  0  0  0  4
## B    0  0  0  0  1  1  2  1  1  1  1  2  1  0  0 11
## C    0  0  0  0  0  0  0  0  0  0  0  0  0  1  1  2
## Sum  1  1  1  1  1  1  2  1  1  1  1  2  1  1  1 17
```


Basics about charts

Principles for good charts according to E. Tufte:

(The Visual Display of Quantitative Information. Cheshire/ Connecticut: Graphics Press, 1983)

- „Graphical excellence is that which gives to the viewer the greatest number of ideas in the shortest time with the least ink in the smallest space.“
- Data-ink ratio = „proportion of a graphic’s ink devoted to the non-redundant display of data-information“ (kein chartjunk!)
- „Graphical excellence is often found in simplicity of design and complexity of data.“

- after Müller-Scheeßel




Plot [1]

Basic drawing function of R:

```
plot(muensingen$Length)
```


options:

- p – points (default)
- l – solid line
- b – line with points for the values
- c – line with gaps for the values
- o – solid line with points for the values
- h – vertical lines up to the values
- s – stepped line from value to value
- n – empty coordinate system




Plot [2]

```
plot(muensingen$Length, type="b")
```



Intelligent system: automatic determination of variable type, drawing of the appropriate chart


```
plot(as.factor(muensingen$fibula_scheme))
```



Plot [3]

Enhancing the plot with optional components & Text

```
plot(muensingen$Length, muensingen$FL,  
     xlim=c(0, 140), # limits of the x axis  
     ylim = c(0, 100), # limits of the y axis  
     xlab = "Fibula Length", # label of the y axis  
     ylab = "Foot Length", # label of the x axis  
     main = "Fibula total length vs. Foot Length", # main title  
     sub="example plot" # subtitle  
     )
```



Plot [4]

Plot do a lot for you:

- Opens a window for display
- Determines the optimal size of the frame of reference
- Draws the coordinate system
- Draws the values

Gives a „handle“ back for further additions to the plot, e.g.:


- lines – additional lines to an existing plot
- points – additional points to an existing plot
- abline – additional special lines to an existing plot
- text – additional text on chosen position to an existing plot

Additional possibilities for “decorations”: ? par

Plot [5]

Add additional elements: Drawing lines

```
abline(v = mean(muensingen$Length), col = "red")           # draw a red vertical line  
abline(h = mean(muensingen$FL), col = "green")             # draw a green vertical line  
abline(lm(FL~Length, data = muensingen), col = "blue")    # draw a blue diagonal line
```



Export the graphics

With the GUI:

Export → Save as...

With the commando line: As vector file

```
dev.copy2eps(file="test.eps")  
dev.copy2pdf(file="test.pdf")
```

```
savePlot(filename="test.tif", type="tiff")
```

Possible are “png”, “jpeg”, “tiff”, “bmp” SavePlot can save sometimes also vector files (dependent on operation system and installation)

Pie chart [1]

The classical one – but also with R not much better...

Used to display proportions, suitable for nominal data

$$a_i = \frac{n_i}{N} * 360^\circ$$

Disadvantages:


- Color selection can influence the perception (red is seen larger than gray)
- Small differences are not easy visible

totally No-Go: 3d-pies!!!

Pie chart [2]

I eat pie...

Ich esse Torte ...



The pieces »viel zu wenig«, »etwas zu wenig« und »gerade richtig« have exactly the same size, the piece »viel zu viel« is a bit smaller. source: <http://www.lrz-muenchen.de/~wlm>


Pie chart [3]

Data are a vector of counts

```
table(muensingen$fibula_scheme)
```

```
##  
##  A  B  C  
##  4 11  2
```


```
pie(table(muensingen$fibula_scheme))
```



Color palette:

The standard palette is pastel, if you prefer another:


```
pie(table(muensingen$fibula_scheme),  
     col=c("red", "green", "blue"))
```




Bar plot [1]

Generally the better alternative... Bar plots are suitable for display of proportions as well as for absolute data. They can be used for every level of measurement.

```
barplot(table(muensingen$fibula_scheme))
```




```
barplot(muensingen$Length)
```



Bar plot [2]

With names:


```
par(las=2) # turn labels 90°  
barplot(muensingen$Length, # plot fibulae length  
        names.arg=muensingen$Grave) # with names of the graves  
title("Fibulae length") # add title
```



Bar plot [3]

Horizontal:

```
par(las=1) # turn labels back again
barplot(table(muensingen$fibula_scheme), # Plot counts fibulae scheme
        horiz=T, # horizontal
        cex.names=2) # make the labels bigger
```




Bar plot [4]

Display of counts


```
my_new_table <- table(muensingen$fibula_scheme,  
                      muensingen$Coils)  
my_new_table
```

```
##  
##      3 4 6 10 12  
## A 1 3 0 0 0  
## B 0 3 6 1 1  
## C 0 1 1 0 0
```

```
barplot(my_new_table)
```



```
barplot(my_new_table, beside=T, legend.text=T)
```




Bar Plot [5]

Display of proportions

```
table.prop<-prop.table(my_new_table,2)
table.prop
```


```
##
##           3           4           6           10           12
## A 1.0000000 0.4285714 0.0000000 0.0000000 0.0000000
## B 0.0000000 0.4285714 0.8571429 1.0000000 1.0000000
## C 0.0000000 0.1428571 0.1428571 0.0000000 0.0000000
```

```
barplot(table.prop)
```



```
tmp<-barplot(table.prop,
             legend.text=T, # add a legend
             col=rainbow(3) # make it more colorful
            )

# add a title
title("ratio of fibulae schemes \n by number of coils",
      outer=TRUE, # outside the plot area
      line=- 3) # on line -3 above
```




Bar Plot [6]

Problems with bar plots – and also with many other charts


Percent vs. count: percents often distort the relations

```
par(mfrow=c(2,1))
barplot(my_new_table,beside=T)
barplot(table.prop,beside=T)
```



Scales: the chosen limits of the axes can distort the relations

```
par(mfrow=c(1,2))
barplot(muensingen$Length[1:2],xpd=F,ylim=c(45,55))
barplot(muensingen$Length[1:2],xpd=F)
```



```
par(mfrow=c(1,1))
```

Box-plot (Box-and-Whiskers-Plot)


One of the best (my precious)!

Used to display the distribution of values in a data vector of metrical (interval, ratio) scale

1 2 3 4 5 6 7 8 9
____|____|____|____


- thick line: mean
- Box: the inner both quantiles
- Whisker: last value < than 1.5 times the distance of the inner quantile

```
boxplot(1:9)
```




Box Plot [2]

```
boxplot(muensingen$Length)
```




```
boxplot(muensingen$Length ~  
        muensingen$fibula_scheme)
```



Box Plot [3]

More beautiful:

```
par(las=1)
boxplot(Length ~ fibula_scheme,
        data = muensingen,
        main = "Length by type",
        col="grey",
        xlab="fibulae scheme",
        ylab="length"
        )
```




Scatterplot [1]

For 2 variables

Used to display a variable in relation to another one. Generally for all scales suitable, but for nominal and ordinal scale other charts are often better.


```
plot(muensingen$Length, muensingen$FL)  
abline(  
  lm(muensingen$FL~muensingen$Length),  
  col="red")
```




Scatterplot [2]

Call additional libraries:

```
library(car) # library for regression analysis
scatterplot(FL ~ Length, data = muensingen)
```




```
library(ggplot2) # advanced plots library
b<- ggplot(muensingen,aes(x=Length,y=FL))
graph<-b + geom_point()
show(graph)
```




Histogramm [1]

Used for classified display of distributions Data reduction vs. precision: Display of count values of classes of values

```
hist(muensingen$Length)
```




```
hist(muensingen$Length, labels = T)
```




Histogramm [2]

Custom breaks of classes

```
hist(muensingen$Length,  
     labels = T)
```




```
hist(muensingen$Length,  
     labels = T,  
     breaks = 10)
```



Histogramm [3]

More beautiful

```
hist(muensingen$Length,breaks=10,  
     labels=T,  
     col="red",  
     xlab="Length",  
     main="Histogram Fibulae Length")
```



Disadvantages:

- Data reduction vs. precision → loss of information
- Actual display depends strongly on the chosen class width

stem-and-leaf chart

An attempt to overcome the disadvantages of a histogram

Is not very often used. Scales like histograms.

```
stem(muensingen$Length)
```

```
##  
## The decimal point is 2 digit(s) to the right of the |  
##  
## 0 | 34444  
## 0 | 5555566677  
## 1 | 13
```

Advantage:


- Information about the distribution inside the classes and the absolute values are (partly) visible.

kernel smoothing (kernel density estimation)

Another attempt to overcome the disadvantages of a histogram


The distribution of the values is considered and a distribution curve is calculated. Continuous distributions are better displayed, without artificial breaks. Scales like histograms.

```
plot(density(muensingen$Length))
```



Histogram and kernel-density-plot together

```
hist(muensingen$Length, prob=T)  
lines(density(muensingen$Length))
```







Style of charts

Stay honest!

- [dax.csv](#)

Choice of display has a strong influence on the statement.



Style of charts

Stay honest!

Choice of display has a strong influence on the statement.

Clear layout!

Minimise Ratio of ink per shown information!

Use the suitable chart for the data!

Consider nominal-ordinal-interval-ratio scale

Suggestions for charts

| What to display | suitable | not suitable |
|---|--------------------------------|-----------------------------|
| Parts of a whole: few | Pie chart, stacked bar plot | |
| Parts of a whole: few | Stacked bar plot | |
| Multiple answers (ties) | Horizontal bar plot | Pie chart, stacked bar plot |
| Comparison of different values of different variables | Grouped bar plot | |
| Comparison of parts of a whole | Stacked bar plot | |
| Comparison of developments | Line chart | |
| Frequency distribution | Histogram, kernel density plot | |
| Correlation of two variables | scatterplot | |